

The length of a set in the sphere whose polynomial hull contains the origin*

by Franc Forstnerič

Department of Mathematics, University of Wisconsin, Madison, WI 53706, USA

Communicated by Prof. J. Korevaar at the meeting of February 24, 1992

ABSTRACT

Let X be a compact subset of the unit sphere in the complex Euclidean space \mathbb{C}^n such that the origin $0 \in \mathbb{C}^n$ belongs to the polynomial hull of X . Assuming that X is rectifiable in the Hausdorff $(\mathcal{H}^1, 1)$ -sense, it is shown that the length of X is at least 2π .

INTRODUCTION

It is well known that every pure one-dimensional complex variety $A \subset \mathbb{C}^n$ with reasonably “nice” boundary bA satisfies the *isoperimetric inequality*

$$(\text{Length}(bA))^2 \geq 4\pi \text{Area}(A).$$

Moreover, if A contains the origin $0 \in \mathbb{C}^n$ while bA lies outside the ball $B(r) = \{z \in \mathbb{C}^n : |z| < r\}$ of radius r , then $\text{Area}(A) \geq \pi r^2$ and therefore

$$\text{Length}(bA) \geq 2\pi r.$$

(See Chirka [6], p. 180 and p. 195, and Bishop [4].) The same inequalities hold if A is an immersed minimal surface with connected boundary; see Section 7.3 in [5]. The constants in these inequalities are the best possible.

H. Alexander [3] extended the isoperimetric inequality to closed Jordan curves $X \subset \mathbb{C}^n$: If X is not polynomially convex, then the set $A = \hat{X} \setminus X$ is an irreducible one-dimensional subvariety of $\mathbb{C}^n \setminus X$, and the isoperimetric in-

*Supported in part by the Research Council of the Republic of Slovenia.

quality holds when the length and the area are computed using the Hausdorff measures \mathcal{H}^1 resp. \mathcal{H}^2 on \mathbb{C}^n . (For Hausdorff measures see Federer [7, p. 171].) In particular, if X is a closed Jordan curve in the unit sphere $S = \{z \in \mathbb{C}^n : |z| = 1\}$ whose polynomial hull \hat{X} contains the origin $0 \in \mathbb{C}^n$, then we have

$$\mathcal{H}^1(X) \geq 2 \sqrt{\pi} (\mathcal{H}^2(\hat{X}))^{1/2} \geq 2\pi.$$

Recall that the polynomial hull of X is the set

$$\hat{X} = \{z \in \mathbb{C}^n : |f(z)| \leq \sup_X |f| \text{ for all } f \in \mathcal{O}(\mathbb{C}^n)\}.$$

In this article we shall consider the following question that was raised by Stout [8, problem 4.2.2]:

If $X \subset S$ is a compact set in the unit sphere whose polynomially convex hull \hat{X} contains the origin $0 \in \mathbb{C}^n$, must the length of X be at least 2π ?

In [11] Stout proved a weaker inequality $\mathcal{H}^1(X) \geq \sqrt{2}\pi$ that improved the previously known result $\mathcal{H}^1(X) \geq 2$ by Sibony [9]. (The result of Sibony applies also to sets that are not contained in a sphere.)

Here we give a very simple proof of the inequality

$$(*) \quad \mathcal{H}^1(X) \geq 2\pi \text{ when } X \subset S \text{ and } 0 \in \hat{X}$$

for $(\mathcal{H}^1, 1)$ -rectifiable compact sets $X \subset S$. Together with a new result of Mark Lawrence this settles the general case as well.

THE RESULT

DEFINITION. (Federer [7, p.251].)

- (a) A set $X \subset \mathbb{R}^k$ is 1-rectifiable if it is the image of a bounded subset $U \subset \mathbb{R}$ under a Lipschitz continuous mapping $f: U \rightarrow \mathbb{R}^k$.
- (b) X is $(\mathcal{H}^1, 1)$ -rectifiable if $\mathcal{H}^1(X) < \infty$ and almost all of X (with respect to the length \mathcal{H}^1) can be covered by a countable union of 1-rectifiable sets.

Our main result is

THEOREM. *If X is a compact $(\mathcal{H}^1, 1)$ -rectifiable subset of the unit sphere $S \subset \mathbb{C}^n$ such that the origin $0 \in \mathbb{C}^n$ belongs to the polynomial hull \hat{X} , then $\mathcal{H}^1(X) \geq 2\pi$.*

Clearly the result can be stated for the ball $r\mathbb{B}$ of radius r : *If $X \subset rS$ is compact and $(\mathcal{H}^1, 1)$ -rectifiable, and if $0 \in \hat{X}$, then $\mathcal{H}^1(X) \geq 2\pi r$.*

The isoperimetric inequality does not hold in the context of our Theorem. Namely, Alexander constructed in [2] a compact disconnected set $X \subset \mathbb{C}^2$ of finite length whose polynomial hull \hat{X} has infinite area. His set X is not highly pathological, it consists of a countable disjoint union of real-analytic simple closed curves, and $A = \hat{X} \setminus X$ is countable union of analytic subsets of $\mathbb{C}^2 \setminus X$. Obviously X is $(\mathcal{H}^1, 1)$ -rectifiable.

It is unknown whether the isoperimetric inequality holds for compact *connected* sets $X \subset \mathbb{C}^n$ of finite length. Recall that $A = \hat{X} \setminus X$ is then a pure one-dimensional analytic subvariety of $\mathbb{C}^n \setminus X$ (if not empty) according to Alexander [1]. For smooth curves X this had been proved by Stolzenberg [10].

REMARK ADDED TO THE PROOF. It suffices to prove the inequality (*) for sets $X \subset S$ of finite length ($\mathcal{H}^1(X) < \infty$) that are minimal, in the sense that no proper compact subset of X contains 0 in its polynomial hull. Recently Mark Lawrence (private communication) informed me of his new result that such a set is necessarily $(\mathcal{H}^1, 1)$ -rectifiable. Together with our theorem this implies

COROLLARY. *If X is a compact subset of S and $0 \in \hat{X}$ then $\mathcal{H}^1(X) \geq 2\pi$.*

Also, after the completion of the first version of this article, H. Alexander [12] and N. Poletski (private communication) informed me that they had independently proved the estimate (*) by different methods.

PROOF OF THE THEOREM

The result will follow from the following Lemma and an integral geometric formula (Crofton formula) from Federer [7, p. 284].

LEMMA. *If X is a compact subset of $\mathbb{C}^n \setminus \{0\}$ of finite length such that $0 \in \hat{X}$, then almost every real hyperplane $\Sigma \subset \mathbb{C}^n$ passing through the origin intersects X at least at two points.*

Here, “almost every” is meant with respect to the volume measure on the Grassman manifold of real hypersurfaces $0 \in \Sigma \subset \mathbb{C}^n$.

PROOF. Since X has finite length and $0 \notin X$, Fubini’s theorem implies that almost every complex hyperplane $L \subset \mathbb{C}^n$ passing through 0 misses X . Here, “almost every” refers to the volume measure on the Grassman manifold of complex $(n-1)$ -dimensional subspaces of \mathbb{C}^n .

Fix such a hyperplane L , and let Σ be any real hyperplane in \mathbb{C}^n containing L . Then L splits Σ in two open half-planes Σ_+ and Σ_- .

We claim that both Σ_+ and Σ_- intersect X . To see this, let $\pi: \mathbb{C}^n \rightarrow L^\perp$ be the orthogonal projection onto the complex line L^\perp orthogonal to L . If X is disjoint from Σ_+ then $\pi(X) \subset L^\perp$ is disjoint from the real half-line $\pi(\Sigma_+)$, hence $0 \in L^\perp$ lies in the unbounded component of $L^\perp \setminus \pi(X)$. Thus 0 is not in the polynomial hull of $\pi(X)$ in L^\perp and hence 0 is not in the hull of X in \mathbb{C}^n , a contradiction. (This also follows from Oka’s criterion for polynomial convexity [10, p. 263]: since 0 belongs to \hat{X} , we can not move L continuously to infinity without hitting X .)

This shows that every such real hyperplane Σ intersects X in at least two points. Since the remaining set of real hyperplanes through the origin in \mathbb{C}^n has measure zero, the lemma is proved.

We now apply Theorem 3.2.48 in Federer [7, p. 284] as follows. Let B be the intersection of a real hyperplane through the origin in \mathbb{C}^n with the sphere S . By our hypothesis X is $(\mathcal{H}^1, 1)$ rectifiable, it is \mathcal{H}^1 measurable since Hausdorff measures are Borel regular, and $\mathcal{H}^1(X) < \infty$. Clearly B is $m = (2n - 2)$ -rectifiable and \mathcal{H}^m -measurable since it is an m -manifold. Applying Theorem 3.2.48 in [7] to the constant functions $\alpha = 1$ on $A = X$ and $\beta = 1$ on B we get

$$\int_{g \in \mathbb{O}(2n)} \int_{X \cap g(B)} d\mathcal{H}^0 d\theta_{2n}(g) = C \cdot \mathcal{H}^1(X) \cdot \mathcal{H}^m(B)$$

for some constant C depending only on m and n .

Recall that \mathcal{H}^0 is just the counting measure. The lemma implies $\mathcal{H}^0(X \cap g(B)) \geq 2$ for almost all $g \in \mathbb{O}(2n)$ with respect to the volume measure θ_{2n} . Hence we get

$$\mathcal{H}^1(X) \geq 2\theta_{2n}(\mathbb{O}(2n))/C\mathcal{H}^m(B).$$

To calculate the constant on the right hand side we choose X to be the intersection of S with a complex line through the origin, hence $\mathcal{H}^1(X) = 2\pi$. In this case $X \cap g(B)$ contains exactly two points for most $g \in \mathbb{O}(2n)$, hence the inequality above is actually an equality. Thus the value of the right hand side equals $\mathcal{H}^1(X) = 2\pi$.

This completes the proof of the Theorem.

REFERENCES

1. Alexander, H. – Polynomial approximation and hulls in sets of finite linear measure in \mathbb{C}^n . Amer. J. Math. **93**, 65–74 (1971).
2. Alexander, H. – The polynomial hull of a set of finite linear measure in \mathbb{C}^n . J. Analyse Math. **47**, 238–242 (1986).
3. Alexander, H. – The polynomial hull of a rectifiable curve in \mathbb{C}^n . Amer. J. Math. **110**, 629–640 (1988).
4. Bishop, E. – Conditions for analyticity, Michigan Math. J. **11**, 289–304 (1964).
5. Burago, Yu.D. and V.A. Zalgaller – Geometric Inequalities. Grundlehren der Math. Wiss. **285**, Springer, Berlin-Heidelberg-New York (1988).
6. Chirka, E.M. – Complex Analytic Sets. Math. and its Appl. **46**, Kluwer, Dordrecht, (1989). Original in Russian, Nauka, Moskva (1985).
7. Federer, H. – Geometric Measure Theory. Grundlehren der Math. Wiss. **153**, Springer, Berlin-Heidelberg-New York, 1969.
8. Fornaess, J.E. et al. – The several complex variables problem list. Ann Arbor, Sept. **10**, (1991).
9. Sibony, N. – Valeurs au bord de fonctions holomorphes et ensembles polynomialement convexes. Lecture Notes in Math. **578**, Springer, Berlin-Heidelberg-New York (1977).
10. Stolzenberg, G. – Polynomially and rationally convex sets. Acta Math. **109**, 259–289 (1963).
11. Stout, E.L. – Removable sets for holomorphic functions of several complex variables. Publ. Mat. **33**, 345–362 (1989).
12. Alexander, H. – Quasi-isoperimetric inequality for polynomial hulls. Preprint (1992).